
Requirements for successful organic citrus production



- > Pest and disease management
- > Water management
- > Pruning and shade management

88

ORANGES

Integrating citrus production into the farming system



The citrus crop must be combined well with other enterprises so that they complement each other.



African Organic Agriculture Training Manual

Raising citrus rootstocks

1. Select seeds

2. Sow seeds



- Choose seeds from healthy and vigorously growing mother trees.
- Squeeze out the seeds and soak them in warm water at 55 °C for 10 minutes.



 Sow the seeds immediately in seedbeds, mixed with wellprepared compost. 3. Transplant seedlings



 Prick the seedlings into polybags when they have 2 pairs of leaves and a bud.



Budding

1. Select healthy trees from which to obtain bud graft branches




2. Gently slice the bud together with the bark form the bud stick




3. Make a 1¹/₂-inch cut in the shape of a "T" into the bark




4. Insert the bud under the "T" slice and wrap with a budding tape












African Organic Agriculture Training Manual

Agroecological requirements of citrus





Planting of citrus seedlings

1. Dig a hole



4. Water the plant well

2. Add topsoil and compost to the bottom



5. Build an earth wall around the seedling

3. Place the tree into the hole and fill the hole with a mixture of topsoil and compost










African Organic Agriculture Training Manual


Establishing a new organic citrus orchard






African Organic Agriculture Training Manual

Improving an existing citrus orchard



Increasing biodiversity in citrus orchard



Promote beneficial plants such as leguminous crops and beneficial herbs.


Control non-beneficial plants such as aggressive grasses and twinning plants.



African Organic Agriculture Training Manual

Improving soil fertility in citrus orchards


A biologically active soil is the foundation for successful citrus production.





African Organic Agriculture Training Manual

How to promote healthy citrus trees



- Ensure good aeration of the crown through annual pruning.
- Use resistant varieties and disease-free planting material.
- Ensure appropriate planting distance for good aeration of the crown.
- Encourage good soil fertility.



Major citrus pests

| Pest | Preventive measures | Direct control |
|---|--|--|
| Leaf miner (Phyllocnistis citrella) Attacks young leaves and shoots. | Promote diversity to encourage natural enemies around and within the orchard. | Apply neem sprays around young leaves and shoots. |
| Aphids (<i>Toxoptera citricidus</i> and <i>T. aurantii</i>) Important vector of the <i>Citrus</i> <i>tristeza</i> virus | Encourage natural enemies such as ladybird beetles, lacewings, hoverflies and parasitic wasps. | Apply neem sprays around the aphid populations, especially around new shoots and under the leaves. |
| Scales Excrete honeydew causing growth of sooty mould, usually associated with ants. | Encourage natural enemies as scales are attacked by a wide range of parasitic wasps and predators. | Apply botanical sprays to target young stages of the scales. Remove heavily affected branches and leaves. |



Major citrus diseases

| Disease | Preventive measures | Direct control |
|--|--|---|
| Damping off caused by the fungi <i>Rhizoctonia solani</i> , <i>Phytophthora</i> spp. or <i>Pythium</i> spp. | Avoid planting in infected fields. Treat seeds with hot water before planting. Avoid over-watering of seedbeds. Use a mixture of manure and sand as planting media. | None possible |
| Greening disease transmitted by vector insects (<i>Diaphorina citri</i>) or infected plant material | Control the vector using <i>Tamarixia radiata</i> or other natural enemies and predators such as <i>Cycloneda sanguinea</i>. | Control the vector using neem, tephrosia or pyre- thrum botanical mixtures. Remove heavily infected trees. |
| Phaeoramularia fruit and leaf spot caused by the fungus <i>Phaeoramularia angolensis</i> | Apply field hygiene by restricting movement of infected material and removal of all infected materials. | Apply copper based products (Bordeaux mixture or copper oxychloride). |
| Citrus tristeza virus (CTV) transmitted by the aphid <i>Toxoptera citricidus</i> | Avoid sour orange rootstock. Use CTV-free budwood. Control vectors like <i>T. itricidus</i>. | Remove infected trees. |



How to manage water efficiently



- > Increase water retention capacity of the soil by ensuring a high level of organic matter and permanent soil cover. This will reduce the need for irrigation.
- > Use water-saving drip irrigation. This will result in economical use of water.
- > Use good quality irrigation water only.



Postharvest handling of citrus

1. Ensure timely harvesting




2. Wash fruits with a mild detergent and clean water



3. Drip to dry under shade



- Citrus should be picked ripe because it does not continue to ripen after harvest.
- 4. Pack and store well



Avoid high humidity conditions in order to control blue and green mould.



Marketing and certification of citrus production

1. Apply organic production methods

2. Proper post harvest handling

3. Certification



 Use organic fertilizers only.



 Do not use synthetic substances to preserve citrus fruits.



- > Only as a market requirement
- Find markets for citrus and other farm products too.
- Cooperate with other farmers to ensure volumes and continuity.

